



บันทึกข้อความ

ส่วนราชการ คณะวิทยาศาสตร์และเทคโนโลยี

ที่ อว ๐๖๓๐.๐๗/คณ๓

วันที่ ๖ กุมภาพันธ์ ๒๕๖๗


เรื่อง ขอส่งหลักฐานเพื่อเบิกเงินค่าตอบแทนนอกเหนือจากเงินค่าจ้างสำหรับพนักงานมหาวิทยาลัย
ที่ดำรงตำแหน่งทางวิชาการ

เรียน คณบดี

ตามที่ ข้าพเจ้าผู้ช่วยศาสตราจารย์ณัฐพงศ์ วัฒนศิริพงษ์ พนักงานมหาวิทยาลัยสายวิชาการ
ตำแหน่งอาจารย์ประจำหลักสูตรคณิตศาสตร์ประยุกต์ คณะวิทยาศาสตร์และเทคโนโลยี ได้ส่งเอกสารการตีพิมพ์
งานวิจัยที่ได้รับการเผยแพร่ตามเกณฑ์ กพอ.กำหนด เพื่อประกอบการเบิกเงินค่าตอบแทนนอกเหนือจากเงิน
ค่าจ้างสำหรับพนักงานมหาวิทยาลัยที่ดำรงตำแหน่งทางวิชาการ ตามปีงบประมาณ ๒๕๖๖ โดยมีบทความวิจัย
เรื่อง Description of regular and intra-regular ordered semigroups by tripolar fuzzy ideals ในวารสาร
Journal of Mathematics and Computer Science Volume 33 (2024), Issue 3, Pages 290–297,
Published online: 04/01/2024.

ทั้งนี้เพื่อให้เป็นไปตามประกาศมหาวิทยาลัยเรื่องแนวทางปฏิบัติในการเบิกเงินค่าตอบแทน
นอกเหนือจากเงินค่าจ้างสำหรับพนักงานมหาวิทยาลัยสายวิชาการ ที่ดำรงตำแหน่งทางวิชาการ พ.ศ.๒๕๖๒ ข้อ
๖.๓ และข้อ ๗ ข้าพเจ้าจึงขอส่งหลักฐานดังกล่าว เพื่อให้คณะกรรมการบริหารคณะพิจารณากลับกรองผลงาน
ทางวิชาการเบื้องต้น

จึงเรียนมาเพื่อโปรดพิจารณา



(ผู้ช่วยศาสตราจารย์ณัฐพงศ์ วัฒนศิริพงษ์)
อาจารย์ประจำหลักสูตรคณิตศาสตร์ประยุกต์



(อาจารย์ ดร.สุภาวิณี ชันคำ)
ประธานหลักสูตรคณิตศาสตร์ประยุกต์

**แบบกลั่นกรองผลงานทางวิชาการของพนักงานมหาวิทยาลัยสายวิชาการ
ที่ดำรงตำแหน่งทางวิชาการ ของคณะกรรมการวิชาการคณะ**

๑. ชื่อผู้เสนอ ณิชฐพงษ์ วัฒนศิริพงษ์ ตำแหน่งวิชาการ ผู้ช่วยศาสตราจารย์
สังกัดหลักสูตร คณิตศาสตร์ประยุกต์
๒. ชื่อผลงาน Description of regular and intra-regular ordered semigroups by tripolar fuzzy ideals. ในวารสาร Journal of Mathematics and Computer Science Volume 33 (2024), Issue 3, Pages 290–297, Published online: 04/01/2024.
๓. คณะกรรมการวิชาการคณะ ได้ตรวจสอบผลงานตามประกาศมหาวิทยาลัย เรื่อง แนวทางปฏิบัติในการเบิกเงินค่าตอบแทนนอกเหนือจากเงินค่าจ้างสำหรับพนักงานมหาวิทยาลัยสายวิชาการ ที่ดำรงตำแหน่งทางวิชาการ พ.ศ.๒๕๖๒ ข้อ ๖.๓ และข้อ ๗ แล้วพบว่า
- ผลงานทางวิชาการปรากฏในเดือน พฤศจิกายน ปี พ.ศ.๒๕๖๖ และอยู่ในวงรอบปีงบประมาณ ๒๕๖๗ ที่ขอรับการประเมิน (๑ ตุลาคม ๒๕๖๖ – ๓๐ กันยายน ๒๕๖๗)
 - เป็นผลงานที่ไม่ลอกเลียนผู้อื่น
 - ผลงานทางวิชาการที่ผู้เสนอขอรับการประเมินตามเกณฑ์ กพอ. กำหนด
 - บทความวิชาการ ตำรา
 - หนังสือ งานวิจัย/บทความวิจัย
 - ผลงานวิชาการลักษณะอื่นๆ
 - สิ่งประดิษฐ์สร้างสรรค์ ผลงานด้านศิลปะ
 - ผลงานบริการสังคม สารานุกรม
 - งานแปล
 - ผลงานทางวิชาการตามเกณฑ์ที่ กพอ. กำหนด มีการเผยแพร่
 - บทความวิชาการ
 - ในวารสารทางวิชาการ หนังสือรวมบทความที่มีบรรณาธิการ
 - ในหนังสือประมวลผลการประชุมทางวิชาการ (Proceeding)
 - เผยแพร่ตำรา
 - ด้วยการพิมพ์โดยโรงพิมพ์หรือสำนักพิมพ์
 - ด้วยสื่ออิเล็กทรอนิกส์ เช่น ซีดีรอม

เผยแพร่หนังสือ

ด้วยการพิมพ์โดยโรงพิมพ์หรือสำนักพิมพ์

ด้วยสื่ออิเล็กทรอนิกส์ เช่น ซีดีรอม

เผยแพร่บทความวิจัย

ในวารสารทางวิชาการ ในหนังสือรวมบทความที่มีบรรณาธิการ

ในหนังสือประมวลผลการประชุมทางวิชาการ (Proceeding)

สัปดาห์ของผลงาน

ร้อยละ ๑๐๐

ร้อยละ ๖๐

ลงชื่อ

(.....)

ประธานคณะกรรมการวิชาการระดับคณะ

วันที่ / /

หมายเลขโทรศัพท์

แบบแสดงหลักฐานการมีส่วนร่วมในผลงานทางวิชาการ

บทความวิจัย ตำรา หนังสือ งานวิจัย ผลงานทางวิชาการในลักษณะอื่น

ผู้ร่วมงาน จำนวน ๓ คน แต่ละคนมีส่วนร่วมดังนี้ :

ชื่อผู้ร่วมงาน	ปริมาณงานร้อยละและหน้าที่ความรับผิดชอบ
ผศ.ณัฐพงศ์ วัฒนศิริพงษ์ Asst. Prof.Nuttapong Wattanasiripong	ปริมาณงานร้อยละ ๖๐ ความคิดริเริ่ม ตั้งสมมติฐาน รวบรวมข้อมูล และพิสูจน์สมมติฐาน
ผศ.ดร.นฤปนาถ เหล็กโคกสูง Asst. Prof. Dr.Nareupanat Lekkoksung	ปริมาณงานร้อยละ ๑๐ ให้ข้อเสนอแนะในการพิสูจน์สมมติฐาน
รศ. ดร.สมศักดิ์ เหล็กโคกสูง Assoc. Prof. Dr.Somsak Lekkoksung	ปริมาณงานร้อยละ ๓๐ ให้ข้อเสนอแนะในการพิสูจน์สมมติฐาน ปรับปรุงทฤษฎีติดต่อประสานงาน

หมายเหตุ :

Wattanasiripong, N., Lekkoksung, N, & Lekkoksung, S. (2024). Description of regular and intra-regular ordered semigroups by tripolar fuzzy ideals. *Journal of Mathematics and Computer Science*, 33(3), 290-297. <https://dx.doi.org/10.22436/jmcs.033.03.07>

ลงชื่อ 
(ผู้ช่วยศาสตราจารย์ณัฐพงศ์ วัฒนศิริพงษ์)

ลงชื่อ 
(ผู้ช่วยศาสตราจารย์ ดร.นฤปนาถ เหล็กโคกสูง)

ลงชื่อ 
(รองศาสตราจารย์ ดร.สมศักดิ์ เหล็กโคกสูง)

JMCS

Journal of Mathematics and Computer Science

Publisher **Home** About Us Announcements Editorial Board Members Journal Info ▾

Sign in ▾ Register

All Fields

Paper Title

Author

Search



Journal of Mathematics and Computer Science (JMCS)

Online: ISSN 2008-949X

Journal of Mathematics and Computer Science-JMCS (J MATH COMPUT SCI-JM) is an international research journal devoted to the publication of significant articles in all areas and sub-areas of mathematics and computer science with applications. High visibility: Indexed in the [Emerging Sources Citation Index - ESCI \(Web of Science\)](#) and [Scopus](#) from Vol. 18 (2018). Submitted papers must be correct, original, nontrivial, a... [Read more](#)

Scopus



Volumes

- Volume 33 (2024)
- Volume 33, Issue 4 pp. 326-430 (2024)

Volume 33, Issue 4, 326-430

Articles: 1 - 8

Certain properties on Bell-based Apostol-type Frobenius-Genocchi polynomials of complex variable

Author(s): A. Al edamat, W. A. Khan, C. S. Ryoo

Pages: 326-338

This journal is a member of, and subscribes to the principles of, the Committee on Publication Ethics (COPE) publicationethics.org



Volumes

- Volume 33 (2024)

- Volume 33, Issue 4
pp. 326-430 (2024)

- Volume 33, Issue 3 pp. 217-325 (2024)

- Volume 33, Issue 2
pp. 124-216 (2024)

- Volume 33, Issue 1
pp. 1-123 (2024)

+ Volume 32 (2024)

+ Volume 31 (2023)

+ Volume 30 (2023)

+ Volume 29 (2023)

+ Volume 28 (2023)

+ Volume 27 (2022)

+ Volume 26 (2022)

+ Volume 25 (2022)

+ Volume 24 (2022)

+ Volume 23 (2021)

+ Volume 22 (2021)

+ Volume 21 (2020)

+ Volume 20 (2020)

+ Volume 19 (2019)

+ Volume 18 (2018)

Volume 33, Issue 3, 217-325

Articles: 1 - 8

Superlinear distributed deviating arguments to study second-order neutral differential equations

Author(s): M. Vijayakumar, S. K. Thamilvanan, B. Sudha, Sh. S. Santra, D. Baleanu
Pages: 217–224

Application of Laplace transform to solve fractional integro-differential equations

Author(s): Gunasekar Gunasekar, P. Raghavendran, Sh. S. Santra, D. Majumder, D. Baleanu, H. Balasundaram
Pages: 225–237

Analyzing convex univalent functions on semi-infinite strip domains

Author(s): V. S. Masih, R. Saadeh, M. Fardi, A. Qazza
Pages: 238–249

Soft algebraic structures embedded with soft members and soft elements: an abstract approach

Author(s): M. Saeed, I. Shafique, A. U. Rahman, S. El-Morsy
Pages: 250–263

An additive-cubic functional equation in a Banach space

Author(s): S. Paokanta, C. Park, N. Jun-on, R. Suparatulorn
Pages: 264–274

A novel decision-making technique based on T-rough bipolar fuzzy sets

Author(s): N. Malik, M. Shabir, T. M. Al-shami, R. Gul, M. Arar
Pages: 275–289

Description of regular and intra-regular ordered semigroups by tripolar fuzzy ideals

Author(s): N. Wattanasiripong, N. Lekkoksung, S. Lekkoksung
Pages: 290–297

Effect of ACE2 receptor and CTL response on within-host dynamics of SARS-CoV-2 infection

Author(s): A. M. Elaiw, A. S. Alsulami, A. D. Hobiny
Pages: 298–325

This journal is a member of, and subscribes to the principles of, the Committee on Publication Ethics (COPE) publicationethics.org



Announcements

JIF WoS: 2.5 (2022)

June 29, 2023

The 2022 Journal Impact Factors have been published by Clarivate Analytics. Our IF (WoS) is now 2.5 for 2022. Our JIF (WoS) is now 1.47. Q1 [Read more](#)

CiteScore 2022: 2.8 Q1

June 05, 2023

CiteScore 2022: 2.8 Q1 [Read more](#)

Acceptance Rate: 0.174

May 25, 2023

Acceptance Rate: 0.174 [Read more](#)

Most Downloaded Articles

Most Visited Articles

Numerical Solutions for Linear Fractional Differential Equations of Order $1 < \alpha < 2$ Using Finite Difference Method (ffdm)

Author(s): Ramzi B. Albadarneh, Iqbal M. Batiha, Mohammad Zurigat
Volume: 16, Issue: 1, Pages: 103-111

SJR Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

Home Journal Rankings Country Rankings Viz Tools Help About Us

EXPLORE, SHOP, AND EMBRACE
THE DRAGON YEAR!

Fly direct from Bangkok (Suvarnabhumi)
✕ Seoul • Tokyo • Osaka
Sapporo • Shanghai
Sydney

FLYER FARE FROM THB **3,290**

Booking Period: 29 Jan 2024 - 4 Feb 2024
Travel Period: 29 Jan 2024 - 31 Oct 2024
*T&C apply

Premium flatbed from THB14,990

Explore, Shop, and Embrace the Dragon Year! All-in one way from THB 3,290

Thai AirAsia X [Book Now >](#)

International Journal of Applied Mathematics and Computer Science

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
Germany	Computer Science └ Computer Science (miscellaneous)	Walter de Gruyter GmbH	50

PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	1641876X	2001-2002, 2006-2022	Homepage How to publish in this journal J.Korbicz@issi.uz.zgora.pl




← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool.**



JMCS

Journal of Mathematics and Computer Science

[Publisher](#)[Home](#)[About Us](#)[Announcements](#)[Editorial Board Members](#)[Journal Info](#) ▾[Sign in](#) ▾[Register](#)

Editorial Board Members

Chief Editor

Martin Bohner - Mathematical Modeling (Department of Mathematics and Statistics, Missouri University of Science and Technology, Rolla, Missouri 65409-0020, USA) [Website](#) [Email](#)

Honorary Chairman

Witold Pedrycz - Computer Sciences (University of Alberta Department of Electrical and Computer Engineering, 9107 - 116 Street, Edmonton, T6G 2V4, Alberta, Canada) [Website](#) [Email](#)

Hari M. Srivastava - (University of Victoria, Victoria, British Columbia V8W 3R4, Canada) [Website](#) [Email](#)

Managing Editor

Tofigh Allahviranloo - Mathematical Modeling (Istinye University, Istanbul, Turkey) [Website](#) [Email](#)

Javad Vahidi - Computer Sciences (Iran University of Science and Technology, Head of Computer Science Department) [Website](#) [Email](#)

Editorial Board

Haydar Akca - Ordinary differential equations, difference equations, numerical analysis, neural networks (Department of Applied Sciences and Mathematics, College of Arts and Sciences, Abu Dhabi University, UAE) [Website](#) [Email](#)

Serkan Araci - q-Series and q-Polynomials, Special functions, Umbral Theory, p-adic analysis, Fractional calculus (Department of Basic Sciences, Faculty of Engineering, Hasan Kalyoncu University, TR-27410, Gaziantep, Turkey) [Website](#) [Email](#)

Abdon Atangana - Applied Mathematics (Institute for Groundwater Studies (IGS) Faculty: Natural and Agricultural Sciences, University of the Free State, Bloemfontein 9300, South Africa) [Website](#) [Email](#)

Dumitru Baleanu - Nonlinear Analysis (Lebanese American University in Beirut) [Website](#) [Email](#)

Martin Bohner - Differential equations, difference equations, dynamic equations on time scales, applications (Department of Mathematics and Statistics Missouri University of Science and Technology Rolla, Missouri 65409-0020, USA) [Website](#) [Email](#)

Dragan Djurcic - Real functions, sequences, series, summability (Kragujevac University) [Website](#) [Email](#)

Rahmat Ellahi - Differential Equations, Mathematical Modeling, Nano Materials, Heat and Mass Transfer Analysis, Blood Flow, Peristaltic Flow, Numerical Simulation, Non-linear Analysis, MHD, Porous Media, Slip, Fluid Mechanics (Department of Mathematics and Statistics, International Islamic University, Islamabad 44000, Pakistan) [Website](#) [Email](#)

Shapour Heidarkhani - Nonlinear Analysis, Differential Equations, Difference Equations, Fractional Differential Equations (Department of Mathematics, Faculty of Sciences, Razi University, 67149 Kermanshah, Iran) [Website](#) [Email](#)

Snezhana Hristova - Ordinary differential equations, fractional differential equations, difference equations, integral equations (Faculty of Mathematics and Informatics, University of Plovdiv, Plovdiv, Bulgaria) [Website](#) [Email](#)

Irena Jadlovská - Differential equations, difference equations, dynamic equations, oscillation theory (Slovak Academy of Sciences, Slovakia) [Website](#) [Email](#)

Hossein Jafari - Numerical analysis, fractional differential equations (Department of Mathematical Sciences, University of South Africa (UNISA), Pretoria 0003, South Africa) [Website](#) [Email](#)

Tongxing Li - Nonlinear analysis, differential and difference equations, oscillation theory (Qingdao Technological University, China) [Website](#) [Email](#)

Jesus Rodríguez Lopez - General topology, geometry, computer science (Polytechnical University of Valencia, Spain) [Website](#) [Email](#)

Rodica Luca - Ordinary and partial differential equations, difference and functional equations, integral equations, positive solutions (Technical University Gh. Asachi, Romania) [Website](#) [Email](#)

Francisco Gallego Lupianez - Topology and fuzzy mathematics (Dept. of Mathematics, Universidad Complutense de Madrid, 28040 Madrid, SPAIN) [Website](#) [Email](#)

Show

Agnieszka Malinowska - Calculus of variations, optimization, difference and differential equations, fractional calculus (Bialystok University of Technology, Poland) [Website](#) [Email](#)

Eberhard Malkowsky - Summability, topology, operator theory, functional analysis (Department of Mathematics, State University of Novi Pazar, Vuka Karadžića bb, 36300 Novi Pazar, Serbia) [Website](#) [Email](#)

Jaqueline Mesquita - dynamic equations on time scales, impulsive differential equations, functional differential equations (Universidade de Brasília, Departamento de Matemática, 70910-900 Brasília, DF, Brazil) [Website](#) [Email](#)

Pedro Tirado Pelaez - Fixed point (Polytechnical University of Valencia, Spain) [Website](#) [Email](#)

Sandra Pinelas - Ordinary differential equations, difference equations, integral equations (Academia Militar Lisboa, Portugal) [Website](#) [Email](#)

Michael Rassias - Number theory, real functions, special functions, difference equations (Department of Mathematics and Engineering Sciences, Hellenic Military Academy, Athens, Greece) [Website](#) [Email](#)

Pradip Roul - Numerical Analysis (Visvesvaraya National Institute of Technology (VNIT)-Nagpur, India) [Website](#) [Email](#)

Ghadir Sadeghi - Functional analysis, operator theory, (noncommutative) metric geometry, (noncommutative) probability spaces, difference and functional equations, fixed point theory (Hakim Sabzevari University, Iran) [Website](#) [Email](#)

Sahra Sedigh Sarvestani - Dependable networks and systems, cyber-physical systems, quantitative modeling and prediction for complex networked systems, critical infrastructure (Missouri S&T, USA) [Website](#) [Email](#)

Sabrina Streipert - Math Biology, Differential Equations, Discrete Mathematics, Time Scales, Population Dynamics, Epidemiology. (University of Pittsburgh) [Website](#) [Email](#)

Gnana Bhaskar Tenali - Ordinary and partial differential equations, fixed point theory (Florida Institute of Technology, USA) [Website](#) [Email](#)

Oscar Valero - Computer science, general topology, functional analysis (Balearic Islands University, Department of Mathematics and Computer Science, Palma, 07122, Balears, Spain) [Website](#) [Email](#)

Calogero Vetro - General topology, operator theory, fixed point theory (University of Palermo, Department of Mathematics and Computer Science, Via Archirafi 34, Palermo 90123, Italy) [Website](#) [Email](#)

Peiguang Wang - Nonlinear analysis, differential and difference equations, stability theory (School of Mathematics and Information Science, Hebei University, The May 4th Street, 071000, China) [Website](#) [Email](#)

Language Editors

J. Math. and Computer Sci.-Jmcs - [Email](#)

Alireza Yeganehparast - Latex and Language Editor (Laval University, Quebec, Canada) [Email](#)

JMCS

Copyright © 2010 - 2024 JMCS. All rights reserved.

Show

Managing Editor

Tofigh Allahviranloo - Mathematical Modeling (Istinye University, Istanbul, Turkey) [Website](#) [Email](#)

Javad Vahidi - Computer Sciences (Iran University of Science and Technology, Head of Computer Science Department) [Website](#) [Email](#)

Editorial Board

Haydar Akca - Ordinary differential equations, difference equations, numerical analysis, neural networks (Department of Applied Sciences and Mathematics, College of Arts and Sciences, Abu Dhabi University, UAE) [Website](#) [Email](#)

Serkan Araci - q-Series and q-Polynomials, Special functions, Umbral Theory, p-adic analysis, Fractional calculus (Department of Basic Sciences, Faculty of Engineering, Hasan Kalyoncu University, TR-27410, Gaziantep, Turkey) [Website](#) [Email](#)

Abdon Atangana - Applied Mathematics (Institute for Groundwater Studies (IGS) Faculty: Natural and Agricultural Sciences, University of the Free State, Bloemfontein 9300, South Africa) [Website](#) [Email](#)

Dumitru Baleanu - Nonlinear Analysis (Lebanese American University in Beirut) [Website](#) [Email](#)

Martin Bohner - Differential equations, difference equations, dynamic equations on time scales, applications (Department of Mathematics and Statistics Missouri University of Science and Technology Rolla, Missouri 65409-0020, USA) [Website](#) [Email](#)

Dragan Djurcic - Real functions, sequences, series, summability (Kragujevac University) [Website](#) [Email](#)

Rahmat Ellahi - Differential Equations, Mathematical Modeling, Nano Materials, Heat and Mass Transfer Analysis, Blood Flow, Peristaltic Flow, Numerical Simulation, Non-linear Analysis, MHD, Porous Media, Slip, Fluid Mechanics (Department of Mathematics and Statistics, International Islamic University, Islamabad 44000, Pakistan) [Website](#) [Email](#)

Shapour Heidarkhani - Nonlinear Analysis, Differential Equations, Difference Equations, Fractional Differential Equations (Department of Mathematics, Faculty of Sciences, Razi University, 67149 Kermanshah, Iran) [Website](#) [Email](#)

Snezhana Hristova - Ordinary differential equations, fractional differential equations, difference equations, integral equations (Faculty of Mathematics and Informatics, University of Plovdiv, Plovdiv, Bulgaria) [Website](#) [Email](#)

Irena Jadlovská - Differential equations, difference equations, dynamic equations, oscillation theory (Slovak Academy of Sciences, Slovakia) [Website](#) [Email](#)

Hossein Jafari - Numerical analysis, fractional differential equations (Department of Mathematical Sciences, University of South Africa (UNISA), Pretoria 0003, South Africa) [Website](#) [Email](#)


Tongxing Li - Nonlinear analysis, differential and difference equations, oscillation theory (Qingdao Technological University, China) [Website](#) [Email](#)

Jesus Rodríguez Lopez - General topology, geometry, computer science (Polytechnical University of Valencia, Spain) [Website](#) [Email](#)

Rodica Luca - Ordinary and partial differential equations, difference and functional equations, integral equations, positive solutions (Technical University Gh. Asachi, Romania) [Website](#) [Email](#)

Francisco Gallego Lupianez - Topology and fuzzy mathematics (Dept. of Mathematics, Universidad Complutense de Madrid, 28040 Madrid, SPAIN) [Website](#) [Email](#)

Show



Description of regular and intra-regular ordered semigroups by tripolar fuzzy ideals



Nuttapong Wattanasiripong^a, Nareupanat Lekkoksung^b, Somsak Lekkoksung^{b,*}

^a*Division of Applied Mathematics, Faculty of Science and Technology, Valaya Alongkorn Rajabhat University under the Royal Patronage, Pathumthani 13180, Thailand.*

^b*Division of Mathematics, Faculty of Engineering, Rajamangala University of Technology Isan, Khon Kaen Campus, Khon Kaen 40000, Thailand.*

Abstract

A mathematical notion dealing with tripolar information is called tripolar fuzzy sets. It was first used to study the properties of Γ -semigroups. In 2022, ordered semigroups were the first considered by tripolar fuzzy sets. It was established that in regular and intra-regular ordered semigroups, the concepts of tripolar fuzzy interior ideals and tripolar ideals are equivalent. The primary goal of this study is to employ tripolar fuzzy left and right ideals to characterize regular and intra-regular ordered semigroups. Additionally, a connection between tripolar fuzzy left (resp., right) ideals and left (resp., right) ideals in ordered semigroups is given.

Keywords: Ordered semigroup, tripolar fuzzy left ideal, tripolar fuzzy right ideal, regularity.

2020 MSC: 08A72, 20M12.

©2024 All rights reserved.

1. Introduction

The idea of fuzzy sets is a mathematical concept that deals with uncertainty in real-world problems. Zadeh [10] invented fuzzy sets for the first time in 1965. Fuzzy sets can be used to study topics in computer science, artificial intelligence, control engineering, robotics, and automata theory. Although fuzzy sets can be used in many scientific investigations, they are limited to several decision-related issues. Mathematical objects called intuitionistic and bipolar fuzzy sets take care of issues that fuzzy sets cannot handle in some circumstances. Atanassov [1] introduced fuzzy sets with intuitive properties. This idea can display both the degree and non-degree of memberships, which helps with ambiguity. Therefore, it can be used to study decision-making problems. Another notion that can be dealt with decision-making difficulties is bipolar fuzzy sets. In 1994, Zhang [11] discussed this concept. There was a discussion of the differences between intuitionistic and bipolar fuzzy sets in 2001 (see [6]).

As a generalization of intuitionistic and bipolar fuzzy sets, Rao [7] presented the idea of tripolar fuzzy sets. Intuitionistic and bipolar fuzzy sets are combined to create this concept. One can use tripolar fuzzy

*Corresponding author

Email addresses: nuttapong@vru.ac.th (Nuttapong Wattanasiripong), nareupanat.le@rmuti.ac.th (Nareupanat Lekkoksung), lekkoksung_somsak@hotmail.com (Somsak Lekkoksung)

doi: [10.22436/jmcs.033.03.07](https://doi.org/10.22436/jmcs.033.03.07)

Received: 2022-07-30 Revised: 2023-12-11 Accepted: 2023-12-14

sets to examine a scenario with tripolar information. When tripolar fuzzy sets were initially introduced, this notion was used for the first time to explore the structural characteristics of Γ -semigroups (see [7]). Later, tripolar fuzzy sets were used in [8] to evaluate the properties of Γ -semirings. The current authors used tripolar fuzzy sets to examine ideals in ordered semigroups. Wattanasiripong et al. [9] presented many types of tripolar fuzzy ideals in ordered semigroups. They demonstrated that in regular and intra-regular ordered semigroups, tripolar fuzzy interior and tripolar (two-sided) ideals coincide.

By the above motivation, we aim to characterize regular and intra-regular ordered semigroups in which any tripolar fuzzy interior and (two-sided) ideal are the same. Before that, we discuss the tripolar fuzzy left and right ideals: the tools used to characterize such classes of ordered semigroups. Additionally, we represent left and right ideals in ordered semigroups using tripolar fuzzy left and right ideals. The paper is organized as follows. Section 2 provides all the necessary basic knowledge used in this paper. Moreover, Some known results are also presented. In our primary section, Section 3, we present the main objectives of our paper. We summarize our detail in Section 4.

2. Preliminaries

This section recalls some basic terminologies of ordered semigroups and tripolar fuzzy sets, which will be used in the paper.

The algebraic system $\langle S; \cdot, \leq \rangle$ is said to be an *ordered semigroup* if $\langle S; \cdot \rangle$ is a semigroup and $\langle S; \leq \rangle$ is a partially ordered set such that $a \cdot c \leq b \cdot c$ and $c \cdot a \leq c \cdot b$ whenever $a \leq b$ for any $a, b, c \in S$. We denote an ordered semigroup $\langle S; \cdot, \leq \rangle$ by the bold letter \mathbf{S} of its universe set. Moreover, we denote the product $x \cdot y$ by xy .

Let \mathbf{S} be an ordered semigroup and A, B , and C nonempty subsets of S . We define the sets AB and (C) by $\{ab : a \in A \text{ and } b \in B\}$ and $\{t \in S : t \leq c \text{ for some } c \in C\}$, respectively. A nonempty subset A of S is called a *left (resp., right) ideal* [3] of \mathbf{S} if:

- (1) $SA \subseteq A$ (resp., $AS \subseteq A$);
- (2) $(A) = A$, equivalently, $x \in A$ whenever $x \leq y$ and $y \in A$.

A set A is referred to be an (*two-sided*) *ideal* of \mathbf{S} if it is both a left and a right ideal of \mathbf{S} .

Let \mathbf{S} be an ordered semigroup. Then \mathbf{S} is said to be:

- (1) *regular* [3] if there exists $x \in S$ such that $a \leq axa$ for any $a \in S$;
- (2) *intra-regular* [4] if there exist $x, y \in S$ such that $a \leq xa^2y$ for any $a \in S$.

Several algebraists intensively study the above classes of ordered semigroups. Additionally, these two classes of ordered semigroups can be studied using the previously introduced notions of left and right ideals.

Lemma 2.1 ([2, 5]). *Let \mathbf{S} be an ordered semigroup. Then*

- (1) \mathbf{S} is regular if and only if $R \cap L = (RL)$ for any left ideal L and right ideal R of \mathbf{S} ;
- (2) \mathbf{S} is intra-regular if and only if $L \cap R \subseteq (LR)$ for any left ideal L and right ideal R of \mathbf{S} .

Definition 2.2 ([7]). Let X be a nonempty set. A set f of the form

$$f := \{(x, f^+(x), f^*(x), f^-(x)) : x \in X \text{ and } 0 \leq f^+(x) + f^*(x) \leq 1\},$$

where $f^+ : X \rightarrow [0, 1]$, $f^* : X \rightarrow [0, 1]$, and $f^- : X \rightarrow [-1, 0]$, is called a *tripolar fuzzy set* of X . The membership degree $f^+(x)$ characterizes the extent to which the element x satisfies the property corresponding to tripolar fuzzy set f , $f^*(x)$ characterizes the extent that the element x satisfies the not property (irrelevant) corresponding to tripolar fuzzy set f , and $f^-(x)$ characterizes the extent that the element x satisfies the implicit counter property of tripolar fuzzy set f .

For simplicity, we write for any tripolar fuzzy set $\{(x, f^+(x), f^*(x), f^-(x)) : x \in X \text{ and } 0 \leq f^+(x) + f^*(x) \leq 1\}$ by the notation f . Therefore, the notation $f(x)$ means $(f^+(x), f^*(x), f^-(x))$ for any $x \in X$. Let $A \subseteq X$. A tripolar fuzzy set χ_A of X , where

$$\chi_A^+(x) := \begin{cases} 1, & \text{if } x \in A, \\ 0, & \text{otherwise,} \end{cases} \quad \chi_A^*(x) := \begin{cases} 0, & \text{if } x \in A, \\ 1, & \text{otherwise,} \end{cases} \quad \chi_A^-(x) := \begin{cases} -1, & \text{if } x \in A, \\ 0, & \text{otherwise,} \end{cases}$$

for any $x \in X$, is called *the characteristic tripolar fuzzy set of A in X*. In particular, if $A = X$, then we denote the tripolar fuzzy set χ_A of A in X by 1. On the other hand, if $A = \emptyset$, then we denote χ_A by 0. By the definition of tripolar fuzzy sets of X , we can regard:

- (1) a fuzzy set $f: X \rightarrow [0, 1]$ of X by a tripolar fuzzy set $(f, 1^*, 1^-)$;
- (2) an intuitionistic fuzzy set $(f^+, f^*): X \rightarrow [0, 1] \times [0, 1]$ of X by a tripolar fuzzy set $(f^+, f^*, 1^-)$;
- (3) a bipolar fuzzy set $(f^+, f^-): X \rightarrow [0, 1] \times [-1, 0]$ of X by a tripolar fuzzy set $(f^+, 1^*, f^-)$.

Let $u_1 = (x_1, y_1, z_1), u_2 = (x_2, y_2, z_2) \in [0, 1] \times [0, 1] \times [-1, 0]$ such that $0 \leq x_1 + y_1 \leq 1$ and $0 \leq x_2 + y_2 \leq 1$. Then we denote by $u_1 \leq u_2$ if $x_1 \leq x_2, y_1 \geq y_2$, and $z_1 \geq z_2$. If $u_1 \leq u_2$ and $u_2 \leq u_1$, then we write $u_1 = u_2$. For any two tripolar fuzzy sets f and g of X , we designate by $f \subseteq g$ whenever $f(x) \leq g(x)$ for all $x \in X$. Moreover, we define a tripolar fuzzy set $f \cap g = ((f \cap g)^+, (f \cap g)^*, (f \cap g)^-)$ of X by

$$(f \cap g)^+(x) := \min\{f^+(x), g^+(x)\}, \quad (f \cap g)^*(x) := \max\{f^*(x), g^*(x)\}, \quad (f \cap g)^-(x) := \max\{f^-(x), g^-(x)\},$$

for all $x \in X$.

Remark 2.3. Let χ_A be the characteristic tripolar fuzzy set of A in X , and $x, y \in X$. Suppose that $\chi_A(y) \leq \chi_A(x)$. Then it is not difficult to illustrate that $x \in A$ whenever $y \in A$.

It is possible to examine the structure of ordered semigroups by using the concept of tripolar fuzzy sets. Let S be an ordered semigroup, and $a \in S$. We define $S_a := \{(x, y) \in S \times S : a \leq xy\}$. For any two tripolar fuzzy sets f and g of S , we define a tripolar fuzzy set $f \circ g = ((f \circ g)^+, (f \circ g)^*, (f \circ g)^-)$ of S by

$$(f \circ g)^+(x) := \begin{cases} \bigvee_{(u,v) \in S_x} \{\min\{f^+(u), g^+(v)\}\}, & \text{if } S_x \neq \emptyset, \\ 0, & \text{otherwise,} \end{cases}$$

$$(f \circ g)^*(x) := \begin{cases} \bigwedge_{(u,v) \in S_x} \{\max\{f^*(u), g^*(v)\}\}, & \text{if } S_x \neq \emptyset, \\ 1, & \text{otherwise,} \end{cases}$$

and

$$(f \circ g)^-(x) := \begin{cases} \bigwedge_{(u,v) \in S_x} \{\max\{f^-(u), g^-(v)\}\}, & \text{if } S_x \neq \emptyset, \\ 0, & \text{otherwise,} \end{cases}$$

for all $x \in S$.

We define the notion of tripolar fuzzy ideals of ordered semigroups as follows.

Definition 2.4 ([9]). Let S be an ordered semigroup. A tripolar fuzzy set f of S is called a *tripolar fuzzy left (resp., right) ideal of S* if for any $x, y \in S$:

- (1) $f(xy) \geq f(y)$ (resp., $f(xy) \geq f(x)$);
- (2) $x \leq y$ implies $f(x) \geq f(y)$.

A tripolar fuzzy set of S is called a *tripolar fuzzy (two-sided) ideal of S* if it is a tripolar fuzzy left and a tripolar fuzzy right ideal of S .

Example 2.5. Let $S = \{a, b, c, d\}$. Define a binary operation \cdot on S and an order relation \leq on S as follows.

·	a	b	c	d
a	a	a	a	a
b	a	a	a	a
c	a	b	c	c
d	a	b	d	d

And $\leq := \Delta_S$, where $\Delta_S := \{(x, x) : x \in S\}$. Then $\mathbf{S} := \langle S; \cdot, \leq \rangle$ is an ordered semigroup. We define a tripolar fuzzy sets f and g of S by

x	f ⁺ (x)	f [*] (x)	f ⁻ (x)		x	g ⁺ (x)	g [*] (x)	g ⁻ (x)
a	0.9	0.1	-1		a	0.9	0.1	-0.5
b	0.4	0.4	-0.6	and	b	0.7	0.2	-0.4
c	0.5	0.3	-0.8		c	0.1	0.6	0
d	0.5	0.3	-0.8		d	0.2	0.8	-0.4

By careful calculation, we obtain that f is a tripolar fuzzy left ideal of \mathbf{S} but not a tripolar fuzzy right ideal of \mathbf{S} since there exist $b, c \in S$ such that $f^+(cb) = 0.4 < 0.5 = f^+(c)$. On the other hand, we obtain that g is a tripolar fuzzy right ideal of \mathbf{S} but not a tripolar fuzzy left ideal of \mathbf{S} since there exist $c, d \in S$ such that $g^+(cd) = 0.1 < 0.2 = g^+(d)$.

The illustration above demonstrates the distinction between tripolar fuzzy left ideals and tripolar right ideals in ordered semigroups.

3. Main results

This section provides a connection between (left, right) ideals and tripolar fuzzy (left, right) ideals in ordered semigroups. Later, we characterize tripolar fuzzy (left, right) ideals in ordered semigroups in terms of the operation \circ . Lastly, we apply such characterizations to classify particular classes of ordered semigroups.

Firstly, we show how left ideals relate to tripolar fuzzy sets in ordered semigroups.

Lemma 3.1. *Let \mathbf{S} be an ordered semigroup and A a nonempty subset of S . Then the following statements are equivalent:*

- (1) A is a left ideal of \mathbf{S} ;
- (2) χ_A is a tripolar fuzzy left ideal of \mathbf{S} .

Proof.

(1) \Rightarrow (2) Let $x, y \in S$. If $xy \in A$, then

$$\chi_A^+(xy) = 1 \geq \chi_A^+(y), \quad \chi_A^*(xy) = 0 \leq \chi_A^*(y), \quad \text{and} \quad \chi_A^-(xy) = -1 \leq \chi_A^-(y).$$

If $xy \notin A$, then $y \notin A$. Thus,

$$\chi_A^+(xy) = \chi_A^+(y), \quad \chi_A^*(xy) = \chi_A^*(y), \quad \text{and} \quad \chi_A^-(xy) = \chi_A^-(y).$$

Therefore, $\chi_A(xy) \geq \chi_A(y)$. Suppose that $x \leq y$. If $y \in A$, then $x \in A$. Thus,

$$\chi_A^+(x) = \chi_A^+(y), \quad \chi_A^*(x) = \chi_A^*(y), \quad \text{and} \quad \chi_A^-(x) = \chi_A^-(y).$$

If $y \notin A$, then

$$\chi_A^+(x) \geq 0 = \chi_A^+(y), \quad \chi_A^*(x) \leq 1 = \chi_A^*(y), \quad \text{and} \quad \chi_A^-(x) \leq 0 = \chi_A^-(y).$$

Therefore, $\chi_A(x) \geq \chi_A(y)$. Altogether, we have that χ_A is a tripolar fuzzy left ideal of \mathbf{S} .

(2) \Rightarrow (1) Let $x \in S$ and $y \in A$. Then $\chi_A(xy) \geq \chi_A(y)$. By Remark 2.3, we have that $xy \in A$. Now, let $x, y \in S$ such that $x \leq y$ and $y \in A$. Then $\chi_A(x) \geq \chi_A(y)$. Again, by Remark 2.3, it follows that $x \in A$. Thus, A is a left ideal of S . \square

With some little modifications, we obtain the following result.

Lemma 3.2. *Let S be an ordered semigroup and A a nonempty subset of S . Then the following statements are equivalent:*

- (1) A is a (right) ideal of S ;
- (2) χ_A is a tripolar fuzzy (right) ideal of S .

Lemmas 3.1 and 3.2 show how tripolar fuzzy left (resp., right, two-sided) ideals can be used to describe left (resp., right, two-sided) ideals in ordered semigroups. The following results illustrate the characterizations of tripolar fuzzy ideals using the operation \circ on the set of all tripolar fuzzy sets of S .

Proposition 3.3. *Let S be an ordered semigroup and f a tripolar fuzzy set of S . Then the following statements are equivalent:*

- (1) f is a tripolar fuzzy left ideal of S ;
- (2) f satisfies the following statements:
 - (a) $1 \circ f \subseteq f$;
 - (b) $f(x) \geq f(y)$ whenever $x \leq y$ for all $x, y \in S$.

Proof.

(1) \Rightarrow (2) It is sufficient to show only that $1 \circ f \subseteq f$. Let $a \in S$. If $S_a = \emptyset$, then

$$(1 \circ f)^+(a) = 0 \leq f^+(a), \quad (1 \circ f)^*(a) = 1 \geq f^*(a), \quad \text{and} \quad (1 \circ f)^-(a) = 0 \geq f^-(a).$$

Suppose that $S_a \neq \emptyset$. Then

$$(1 \circ f)^+(a) = \bigvee_{(u,v) \in S_a} \{\min\{1^+(u), f^+(v)\}\} = \bigvee_{(u,v) \in S_a} f^+(v) \leq \bigvee_{(u,v) \in S_a} f^+(uv) \leq \bigvee_{(u,v) \in S_a} f^+(a) = f^+(a),$$

$$(1 \circ f)^*(a) = \bigwedge_{(u,v) \in S_a} \{\max\{1^*(u), f^*(v)\}\} = \bigwedge_{(u,v) \in S_a} f^*(v) \geq \bigwedge_{(u,v) \in S_a} f^*(uv) \geq \bigwedge_{(u,v) \in S_a} f^*(a) = f^*(a),$$

and

$$(1 \circ f)^-(a) = \bigwedge_{(u,v) \in S_a} \{\max\{1^-(u), f^-(v)\}\} = \bigwedge_{(u,v) \in S_a} f^-(v) \geq \bigwedge_{(u,v) \in S_a} f^-(uv) \geq \bigwedge_{(u,v) \in S_a} f^-(a) = f^-(a).$$

This shows that $(1 \circ f)(a) \leq f(a)$. That is, $(1 \circ f) \subseteq f$.

(2) \Rightarrow (1) We need to verify only that $f(xy) \geq f(y)$ for all $x, y \in S$. Let $x, y \in S$. Then

$$f^+(xy) \geq (1 \circ f)^+(xy) = \bigvee_{(u,v) \in S_{xy}} \{\min\{1^+(u), f^+(v)\}\} \geq \min\{1^+(x), f^+(y)\} = f^+(y),$$

$$f^*(xy) \leq (1 \circ f)^*(xy) = \bigwedge_{(u,v) \in S_{xy}} \{\max\{1^*(u), f^*(v)\}\} \leq \max\{1^*(x), f^*(y)\} = f^*(y),$$

and

$$f^-(xy) \leq (1 \circ f)^-(xy) = \bigwedge_{(u,v) \in S_{xy}} \{\max\{1^-(u), f^-(v)\}\} \leq \max\{1^-(x), f^-(y)\} = f^-(y).$$

This shows that $f(xy) \geq f(y)$. That is, f is a tripolar fuzzy left ideal of S . \square

Similarly, we can prove the following propositions.

Proposition 3.4. *Let S be an ordered semigroup and f a tripolar fuzzy set of S . Then the following statements are equivalent:*

- (1) f is a tripolar fuzzy right ideal of S ;
- (2) f satisfies the following statements:
 - (a) $f \circ 1 \subseteq f$;
 - (b) $f(x) \geq f(y)$ whenever $x \leq y$ for all $x, y \in S$.

Proposition 3.5. *Let S be an ordered semigroup and f a tripolar fuzzy set of S . Then the following statements are equivalent:*

- (1) f is a tripolar fuzzy ideal of S ;
- (2) f satisfies the following statements:
 - (a) $f \circ 1 \subseteq f$ and $1 \circ f \subseteq f$;
 - (b) $f(x) \geq f(y)$ whenever $x \leq y$ for all $x, y \in S$.

Later, we apply the notion of tripolar fuzzy ideals in ordered semigroups to characterize three classes of ordered semigroups; regular, intra-regular, and both regular and intra-regular ordered semigroups. We can explain these particular classes of ordered semigroups with the help of the following results.

Lemma 3.6. *Let S be an ordered semigroup, and $A, B \subseteq S$. Then*

- (1) $\chi_A \subseteq \chi_B$ if and only if $A \subseteq B$;
- (2) $\chi_A \circ \chi_B = \chi_{(AB]}$;
- (3) $\chi_A \cap \chi_B = \chi_{A \cap B}$.

Proof. The propositions (1) and (3) are not difficult to prove. Hence, we prove only (2).

(2) Let $x \in S$. It is not difficult to see that if $x \notin (AB]$, that is, there is no $a \in A$ and $b \in B$ such that $x \leq ab$, then

$$\chi_{(AB]}^+(x) = 0 = (\chi_A \circ \chi_B)^+(x), \quad \chi_{(AB]}^*(x) = 1 = (\chi_A \circ \chi_B)^*(x), \quad \text{and} \quad \chi_{(AB]}^-(x) = 0 = (\chi_A \circ \chi_B)^-(x).$$

Suppose that $x \in (AB]$. That is, there exist $a \in A$ and $b \in B$ such that $x \leq ab$. Then

$$\chi_{(AB]}^+(x) = 1 = \min\{\chi_A^+(a), \chi_B^+(b)\} = \bigvee_{(u,v) \in S_x} \{\min\{\chi_A^+(u), \chi_B^+(v)\}\} = (\chi_A \circ \chi_B)^+(x),$$

$$\chi_{(AB]}^*(x) = 0 = \max\{\chi_A^*(a), \chi_B^*(b)\} = \bigwedge_{(u,v) \in S_x} \{\max\{\chi_A^*(u), \chi_B^*(v)\}\} = (\chi_A \circ \chi_B)^*(x),$$

and

$$\chi_{(AB]}^-(x) = -1 = \max\{\chi_A^-(a), \chi_B^-(b)\} = \bigwedge_{(u,v) \in S_x} \{\max\{\chi_A^-(u), \chi_B^-(v)\}\} = (\chi_A \circ \chi_B)^-(x).$$

Therefore, $\chi_A \circ \chi_B = \chi_{(AB]}$. □

Lemma 3.7. *Let S be an ordered semigroup. Then $f \circ g \subseteq f \cap g$ for any tripolar fuzzy left ideal g and tripolar fuzzy right ideal f of S .*

Proof. Let g and f be a tripolar fuzzy left and a tripolar fuzzy right ideal of S , respectively. Given $a \in S$, if $S_a = \emptyset$, then $(f \circ g)(a) = 0(a) \leq (f \cap g)(a)$. Suppose that $S_a \neq \emptyset$. We observe that $f \subseteq 1$ and $g \subseteq 1$. Then, by Propositions 3.3 and 3.4, we have

$$(f \circ g)^+(a) \leq (1 \circ g)^+(a) \leq g^+(a) \quad \text{and} \quad (f \circ g)^+(a) \leq (f \circ 1)^+(a) \leq f^+(a).$$

This implies that $(f \circ g)^+(a) \leq \min\{g^+(a), f^+(a)\}$. Similarly, by Propositions 3.3 and 3.4, we have

$$(f \circ g)^*(a) \geq (1 \circ g)^*(a) \geq g^*(a) \quad \text{and} \quad (f \circ g)^*(a) \geq (f \circ 1)^*(a) \geq f^*(a).$$

This implies that $(f \circ g)^*(a) \geq \max\{g^*(a), f^*(a)\}$. Again, by Propositions 3.3 and 3.4, we have

$$(f \circ g)^-(a) \geq (1 \circ g)^-(a) \geq g^-(a) \quad \text{and} \quad (f \circ g)^-(a) \geq (f \circ 1)^-(a) \geq f^-(a).$$

This implies that $(f \circ g)^-(a) \geq \max\{g^-(a), f^-(a)\}$. Thus, $(f \circ g)(a) \leq (f \cap g)(a)$. Altogether, we obtain our claim. \square

Now, we are ready to characterize regular ordered semigroups using tripolar fuzzy left and right ideals.

Theorem 3.8. *Let \mathbf{S} be an ordered semigroup. Then the following statements are equivalent:*

- (1) \mathbf{S} is regular;
- (2) $f \circ g = f \cap g$ for every tripolar fuzzy left ideal g and tripolar fuzzy right ideal f of \mathbf{S} .

Proof.

(1) \Rightarrow (2) Let g and f be a tripolar fuzzy left and a tripolar fuzzy right ideal of \mathbf{S} , respectively. By Lemma 3.7, we need to illustrate only that $f \cap g \subseteq f \circ g$. Let $a \in \mathbf{S}$. Since \mathbf{S} is regular, there exists $x \in \mathbf{S}$ such that $a \leq axa = a(xa)$. This means that $\mathbf{S}_a \neq \emptyset$. Then

$$(f \circ g)^+(a) = \bigvee_{(u,v) \in \mathbf{S}_a} \{\min\{f^+(u), g^+(v)\}\} \geq \min\{f^+(a), g^+(xa)\} \geq \min\{f^+(a), g^+(a)\},$$

$$(f \circ g)^*(a) = \bigwedge_{(u,v) \in \mathbf{S}_a} \{\max\{f^*(u), g^*(v)\}\} \leq \max\{f^*(a), g^*(xa)\} \leq \max\{f^*(a), g^*(a)\},$$

and

$$(f \circ g)^-(a) = \bigwedge_{(u,v) \in \mathbf{S}_a} \{\max\{f^-(u), g^-(v)\}\} \leq \max\{f^-(a), g^-(xa)\} \leq \max\{f^-(a), g^-(a)\}.$$

This shows that $f \cap g \subseteq f \circ g$. Thus, by Lemma 3.7, $f \circ g = f \cap g$.

(2) \Rightarrow (1) Let L and R be a left and a right ideal of \mathbf{S} , respectively. By Lemmas 3.1 and 3.2, we have that χ_L and χ_R are a tripolar fuzzy left and a tripolar fuzzy right ideal of \mathbf{S} , respectively. By our hypothesis and Lemma 3.6, we have

$$\chi_{R \cap L} = \chi_R \cap \chi_L = \chi_R \circ \chi_L = \chi_{(RL)}.$$

This implies, by Lemma 3.6, that $R \cap L = (RL)$. By Lemma 2.1, \mathbf{S} is regular. \square

In our next result, we describe intra-regular ordered semigroups by tripolar fuzzy left and tripolar fuzzy right ideals of \mathbf{S} .

Theorem 3.9. *Let \mathbf{S} be an ordered semigroup. Then the following statements are equivalent:*

- (1) \mathbf{S} is intra-regular;
- (2) $f \cap g \subseteq f \circ g$ for every tripolar fuzzy left ideal f and tripolar fuzzy right ideal g of \mathbf{S} .

Proof.

(1) \Rightarrow (2) Let f and g be a tripolar fuzzy left and a tripolar fuzzy right ideal of \mathbf{S} , respectively. Let $a \in \mathbf{S}$.

Since \mathbf{S} is intra-regular, there exist $x, y \in S$ such that $a \leq xa^2y = (xa)(ay)$. This means that $S_a \neq \emptyset$. Then

$$(f \circ g)^+(a) = \bigvee_{(u,v) \in S_a} \{\min\{f^+(u), g^+(v)\}\} \geq \min\{f^+(xa), g^+(ay)\} \geq \min\{f^+(a), g^+(a)\},$$

$$(f \circ g)^*(a) = \bigwedge_{(u,v) \in S_a} \{\max\{f^*(u), g^*(v)\}\} \leq \max\{f^*(xa), g^*(ay)\} \leq \max\{f^*(a), g^*(a)\},$$

and

$$(f \circ g)^-(a) = \bigwedge_{(u,v) \in S_a} \{\max\{f^-(u), g^-(v)\}\} \leq \max\{f^-(xa), g^-(ay)\} \leq \max\{f^-(a), g^-(a)\}.$$

This shows that $f \cap g \subseteq f \circ g$.

(2) \Rightarrow (1) Let L and R be a left and a right ideal of \mathbf{S} , respectively. By Lemmas 3.1 and 3.2, we have that χ_L and χ_R are a tripolar fuzzy left and a tripolar fuzzy right ideal of \mathbf{S} , respectively. By our hypothesis and Lemma 3.6, we have

$$\chi_{L \cap R} = \chi_L \cap \chi_R \subseteq \chi_L \circ \chi_R = \chi_{(LR)}.$$

This implies, by Lemma 3.6, that $L \cap R \subseteq (LR)$. By Lemma 2.1, \mathbf{S} is intra-regular. \square

Our final conclusion follows from Theorems 3.8 and 3.9.

Theorem 3.10. *Let \mathbf{S} be an ordered semigroup. Then the following statements are equivalent:*

- (1) \mathbf{S} is both regular and intra-regular;
- (2) $f \cap g \subseteq (g \circ f) \cap (f \circ g)$ for every tripolar fuzzy left ideal f and tripolar fuzzy right ideal g of \mathbf{S} .

4. Conclusion

We concentrate on a generalization of bipolar and intuitionistic fuzzy left (resp., right) ideals in ordered semigroups in this study. We establish a link between tripolar fuzzy left (resp., right) ideals and left (resp., right) ideals. A binary operation defined on the set of all tripolar fuzzy sets of ordered semigroups was used to describe tripolar fuzzy left (resp., right) ideals. Lastly, we use the notion of tripolar fuzzy left (resp., right) ideals to represent regular and intra-regular ordered semigroups. As a result, Lemma 2.1 is generalized by our determinations. Since many algebraic systems can be studied using tripolar fuzzy sets, it is still unknown if the same idea can be utilized to investigate hyperalgebraic systems.

References

- [1] K. T. Atanassov, *Intuitionistic fuzzy sets*, Fuzzy Sets and Systems, **20** (1986), 87–96. 1
- [2] Y. Cao, *Characterizations of regular ordered semigroups by strong quasi-ideals*, Vietnam J. Math., **30** (2002), 239–250. 2.1
- [3] N. Kehayopulu, *On completely regular poe-semigroups*, Math. Japon., **37** (1992), 123–130. 2, 1
- [4] N. Kehayopulu, *On intra-regular ordered semigroups*, Semigroup Forum, **46** (1993), 271–278. 2
- [5] D. M. Lee, S. K. Lee, *On intra-regular ordered semigroups*, Kangweon-Kyungki Math. J., **14** (2006), 95–100. 2.1
- [6] K. M. Lee, K. M. Lee, K. J. Cios, *Comparison of interval-valued fuzzy sets, intuitionistic fuzzy sets, and bipolar-valued fuzzy sets*, Comput. Inf. Technol., (2001), 433–439. 1
- [7] M. M. K. Rao, *Tripolar fuzzy interior ideals of Γ -semigroup*, Ann. Fuzzy Math. Inform., **15** (2018), 199–206. 1, 2.2
- [8] M. M. K. Rao, B. Venkateswarlu, *Tripolar fuzzy interior ideals of a Γ -semiring*. Asia Pac. J. Math., **5** (2018), 192–207. 1
- [9] N. Wattanasiripong, N. Lekkoksung, S. Lekkoksung, *On tripolar fuzzy interior ideals in ordered semigroups*, Int. J. Innov. Comput. Inf. Control, **18** (2022), 1291–1304. 1, 2.4
- [10] L. A. Zadeh, *Fuzzy sets*, Inf. Control, **3** (1965), 338–353. 1
- [11] W. R. Zhang, *Bipolar fuzzy sets and relations: a computational framework for cognitive modeling and multiagent decision analysis*, In: NAFIPS/IFIS/NASA'94. Proceedings of the first international joint conference of the North American fuzzy information processing society biannual conference, The Industrial Fuzzy Control and Intelligence, IEEE, (1994), 305–309. 1